



## King's Research Portal

*Document Version*  
Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Mizerovska, J., Dracinska, H., Arlt, V. M., Schmeiser, H. H., Frei, E., & Stiborova, M. (2009). Oxidation of 3-aminobenzanthrone, a human metabolite of carcinogenic environmental pollutant 3-nitrobenzanthrone, by cytochromes P450-similarity between human and rat enzymes. *Neuroendocrinology Letters*, 30(SUPPL.1), 52 - 59.

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Oxidation of 3-aminobenzanthrone, a human metabolite of carcinogenic  
environmental pollutant 3-nitrobenzanthrone, by cytochromes P450 – similarity  
between human and rat enzymes**

**RNDr Jana Mizerovská<sup>1</sup>, Helena Dračínská PhD<sup>1</sup>, Volker M. Arlt PhD<sup>2</sup>, Heinz H  
Schmeiser PhD<sup>3</sup>, Eva Frei PhD<sup>3</sup>, Prof Marie Stiborová DrSc<sup>1</sup>**

<sup>1</sup>Department of Biochemistry, Faculty of Science, Charles University, Czech Republic

<sup>2</sup>Section of Molecular Carcinogenesis, Institute of Cancer Research, Brookes Lawley  
Building, United Kingdom

<sup>3</sup>German Cancer Research Center, Germany

Corresponding author: Prof. RNDr. Marie Stiborová, DrSc, Department of  
Biochemistry, Faculty of Science, Charles University, Prague, Albertov 2030, 128 40  
Prague 2, Czech Republic, TEL: +420-221951285, fax: +420-221951283, e-mail:  
[stiborov@natur.cuni.cz](mailto:stiborov@natur.cuni.cz)

Running headline: 3-Aminobenzanthrone oxidation by cytochromes P450

**OBJECTIVES:** 3-Aminobenzanthrone (3-ABA) is the main human metabolite of  
carcinogenic environmental pollutant 3-nitrobenzanthrone (3-NBA). Understanding  
which cytochrome P450 (CYP) enzymes are involved in metabolism of this toxicant is  
important in the assessment of individual susceptibility. Characterization of 3-ABA  
metabolites formed by rat hepatic microsomes containing cytochromes P450 (CYPs)

and identification of the major rat and human CYPs participating in this process are aims of this study.

**METHODS:** HPLC with UV detection was employed for the separation and characterization of 3-ABA metabolites. Inducers and inhibitors of CYPs and rat and human recombinant CYPs were used to characterize the enzymes participating in 3-ABA oxidation.

**RESULTS:** Selective CYP inhibitors and hepatic microsomes of rats pre-treated with specific CYP inducers were used to characterize rat liver CYPs metabolizing 3-ABA (measured as consumption of 3-ABA). Kinetics of these reactions catalyzed by rat hepatic microsomes was also evaluated. Based on these studies, we attribute most of 3-ABA metabolism in rat liver to CYP1A and 3A. Among recombinant rat and human CYP enzymes tested in this study, rat CYP3A2 and human CYP3A4/5, followed by CYP1A1 of both organisms were the most effective enzymes converting 3-ABA. Rat hepatic CYP enzymes oxidize 3-ABA up to three metabolites. Two of them were identified to be the products formed by oxidation of 3-ABA on its amino group back to the parent compound from which 3-ABA is generated in organisms, 3-NBA. Namely, *N*-hydroxylation metabolite, *N*-hydroxy-3-ABA and 3-NBA were identified to be these 3-ABA oxidation products. These metabolites are formed by CYPs of a 1A subfamily. Another 3-ABA metabolite, whose structure remains to be characterized, is generated not only by CYP1A but also by other CYP enzymes, predominantly by CYPs of a 3A subfamily.

**CONCLUSIONS:** The results found in this study, the first report on the metabolism of 3-ABA by human and rat CYPs, clearly demonstrate that CYPs of 3A and 1A subfamilies are the major enzymes metabolizing 3-ABA.

50 **KEY WORDS**

51 3-aminobenzanthrone; 3-nitrobenzanthrone, cytochrome P450; metabolism.

52

53 **ABBREVIATIONS & UNITS**

54 3-ABA – 3-aminobenzanthrone

55  $\alpha$ -NF ;  $\alpha$ -naphthoflavone

56  $\beta$ -NF ;  $\beta$ -naphthoflavone

57 cDNA - complementary DNA

58 CYP - cytochrome P450

59 DDTC - diethyldithiocarbamic acid

60 HPLC - high performance liquid chromatography

61  $K_m$  - Michaelis constant

62 MS - microsomes

63 *N*-hydroxy-3-ABA - *N*-hydroxy-3- aminobenzanthrone

64 NADPH - nicotinamidadeninedinucleotide phosphate (reduced)

65 3-NBA – 3-nitrobenzatntrone

66 PB – phenobarbital

67 RP – reverse phase

68 r.t. – retention time

69 UV – ultraviolet

70 VIS - visible

71  $V_{max}$  - maximum reaction rate

72

## INTRODUCTION

The nitroaromatic compound 3-nitrobenzanthrone (3-nitro-7H-benz[de]anthracen-7-one, 3-NBA, Fig. 1) is one of the most potent mutagens and a suspected human carcinogen that is found in diesel exhaust and ambient air pollution (Arlt, 2005, Hansen et al., 2007). We found that 3-NBA is activated to *N*-hydroxy-3-aminobenzanthrone (*N*-hydroxy-3-ABA) by cytosolic and microsomal reductases by simple nitroreduction (Arlt 2005, Arlt et al., 2002, 2003a,b,c, 2005, Stiborova et al., 2006a, 2008, 2009, Svobodova et al. 2007) (Figure 1).

Recently 3-NBA has received much attention due to its extremely high mutagenic potency in the Ames Salmonella assay (Enya et al., 1997; Seidel et al., 2002; Arlt, 2005). 3-NBA is carcinogenic in rats, causing lung tumours after intratracheal instillation, and it is also a suspected human carcinogen (Seidel et al., 2002; Arlt, 2005; Nagy et al., 2005).

The uptake of 3-NBA in humans has been demonstrated by the detection of its metabolite 3-aminobenzanthrone (3-ABA, Figure 1) in urine samples of salt mine workers occupationally exposed to diesel emissions (Seidel et al., 2002). 3-ABA was also the main metabolite of 3-NBA formed in human fetal bronchial cells and rat lung alveolar type II cells (Borlak et al., 2000).

3-ABA was also evaluated to be suitable for coloration of microporous polyethylene films, which are widely used for practical purposes such as separation of liquid mixtures, in particular, as separation membranes in chemical batteries (Grabchev et al., 2002), or an advantageous fluorescent phospholipid membrane label in the form of its *N*-palmitoyl derivative (Sykora et al., 2002). This suggests industrial and/or laboratory utilization of this 3-NBA metabolite, leading to a putative exposure of people. This is a matter of concern, because we have demonstrated the

genotoxicity of both 3-NBA and 3-ABA by the detection of specific DNA adducts formed *in vitro* and *in vivo* (Arlt et al., 2001; 2002; 2003a; b; c; 2004a; c; 2005; 2006b; Bieler et al., 1999; 2005; 2007; Stiborova et al., 2006a; 2008, 2009). Previous work indicated that *N*-hydroxy-3-ABA appears to be the critical intermediate in 3-NBA/ABA-derived DNA adduct formation (Figure 1), which can be further activated by *N,O*-acetyltransferases (NATs) and sulfotransferases (SULTs) (Arlt et al., 2002, 2003a, b, 2005). The predominant DNA adducts formed from 3-NBA and 3-ABA are 2-(2'-deoxyguanosin- $N^2$ -yl)-3-aminobenzanthrone (dG- $N^2$ -ABA) and *N*-(2'-deoxyguanosin-8-yl)-3-aminobenzanthrone (dG-C8-*N*-ABA) and these are most probably responsible for the induction of GC to TA transversion mutations induced by these toxicants (Arlt et al., 2004a; Arlt et al., 2006b; Bieler et al., 2007).

Understanding which enzymes are involved in the metabolism (activation and/or detoxication) of 3-ABA is important in the assessment of susceptibility to this 3-NBA metabolite. Recently, we have found that cytochromes P450 (CYP) 1A1 and 1A2 are essential for 3-ABA oxidative activation in human and rat liver, lung and kidney to reactive species, *N*-hydroxy-3-ABA, forming the same DNA adducts that are formed *in vitro* and *in vivo* in rodents by 3-ABA or 3-NBA (Arlt et al. 2003a, b, c, 2004b, 2005, 2006a, Stiborova et al. 2006a; 2008, 2009) (Figure 1).

In contrast to the enzymes activating 3-ABA to species binding to DNA, those participating in 3-ABA oxidation to other potential metabolites have not been extensively studied so far. Therefore, here we investigated the oxidative metabolism of 3-ABA *in vitro*, in order to characterize the 3-ABA metabolites and to identify CYPs responsible for their formation. Hepatic microsomal systems and recombinant CYP enzymes of rat, the organism found previously to be a suitable model mimicking activation metabolism of 3-ABA in human (Arlt et al., 2004b) were used for such a

study. In addition, rat and human recombinant CYP enzymes were utilized to characterize their participation in 3-ABA oxidation.

## **MATERIAL AND METHODS**

*Synthesis of 3-ABA and N-hydroxy-3-ABA.* 3-ABA and N-hydroxy-3-ABA were synthesized as described by Arlt et al. (2003a) and their authenticity was confirmed by UV spectroscopy, electrospray mass spectra and high field proton NMR spectroscopy.

*Chemicals and enzymes.* Microsomes from rat livers were isolated and characterized for CYP activities as described (Arlt et al., 2004b, Stiborova et al., 2006a,b). Supersomes™, microsomes isolated from insect cells transfected with Baculovirus constructs containing cDNA of one of the following rat CYPs: CYP1A1, 1A2, 2A1, 2A2, 2B1, 2C6, 2C11, 2C12, 2C13, 2D1, 2D2, 2E1, 3A1, 3A2 with cytochrome b<sub>5</sub> and expressing NADPH:CYP reductase and one of the following human CYPs: CYP1A1, 1A2, 2A6, 2B6, 2C8 (with and without cytochrome b<sub>5</sub>), 2C19 (with and without cytochrome b<sub>5</sub>), 2D6, 2E1, 3A4 and 3A5 (with and without cytochrome b<sub>5</sub>), and expressing NADPH:CYP reductase were obtained from Gentest Corp. (USA). NADPH, phenacetine, dimethylsulfoxide (DMSO) were obtained from Sigma Chemical Co (St Louis, MO, USA), ethylacetate, methanol for high performance liquid chromatography (HPLC) super gradient, methanol were obtained from Lachema, (Brno, Czech Republic).

*Animal experiments and preparation of microsomes.* The study was conducted in accordance with the Regulations for the Care and Use of Laboratory Animals (311/1997, Ministry of Agriculture, Czech Republic), which complies with the Declaration of Helsinki. Microsomes from livers of ten male untreated Wistar rats and

148 those from livers of ten male rats pre-treated with  $\beta$ -naphthoflavone ( $\beta$ -NF) and  
149 phenobarbital (PB) were prepared by the procedure described previously (Arlt et al.,  
150 2004b, Stiborova et al., 2002b, 2006b,. Krizkova et al., 2008, Sistkova et al., 2008).  
151 Protein concentrations in the microsomal fractions were assessed using the  
152 bicinchoninic acid protein assay with the bovine serum albumin as a standard  
153 (Weichelman et al., 1988). The concentration of CYP was estimated according to  
154 Omura and Sato (Omura and Sato, 1964) based on absorption of the complex of  
155 reduced CYP with carbon monoxide. Untreated rat liver microsomes contained 0.6  
156 nmol CYP/mg protein. Hepatic microsomes of rats treated with  $\beta$ -NF and PB  
157 contained 1.3 and 1.5 nmol CYP/mg proteins, respectively.

158 *Incubations.* Unless stated otherwise, incubation mixtures used for studying 3-ABA  
159 metabolism by rat hepatic microsomes were prepared as described previously by  
160 Mizerovska et al. (2008). Briefly, incubation mixtures, containing final volume of 500  
161  $\mu$ l, consisted of 100 mM potassium phosphate buffer (pH 7.4), 10 mM NADPH, 0.5  
162 mg of microsomal protein and 5 - 50  $\mu$ M 3-ABA (dissolved in DMSO). The reaction  
163 was initiated by adding 3-ABA. Incubations with rat microsomes were carried out at  
164 37 °C for 5 minutes. Control incubations were carried out either without the  
165 enzymatic system (microsomes) or without NADPH. Incubation mixtures used for  
166 studying 3-ABA metabolism by rat and human recombinant CYP enzymes contained  
167 final volume of 250  $\mu$ l, consisting of 100 mM potassium phosphate buffer (pH 7.4), 10  
168 mM NADPH, 1  $\mu$ M CYP in Supersomes<sup>TM</sup> and 20  $\mu$ M 3-ABA (dissolved in DMSO).  
169 All incubations were carried out at 37 °C for 20 min. In the case of investigation of the  
170 time dependence of 3-ABA oxidation, reaction mixtures were incubated at 37°C for 0  
171 - 120 minutes. Control incubations were carried out either without the CYP enzymes  
172 (Supersomes<sup>TM</sup>) or without NADPH. Then, 2.5 or 5  $\mu$ l of 1 mM phenacetine in



methanol was added as an internal standard, and 3-ABA and its metabolites were extracted twice with ethyl acetate (2 x 1.5 ml). The extracts were evaporated to dryness; residues dissolved in 30 µl of methanol and subjected to reverse-phase (RP)-HPLC to evaluate the amounts of residual 3-ABA and its metabolites. The 3-ABA metabolites were separated from 3-ABA by HPLC with UV detection and characterized by mass spectrometry and co-chromatography with synthetic standards. The following chemicals were used in the inhibition studies of the 3-ABA oxidation by hepatic microsomes:  $\alpha$ -naphthoflavone ( $\alpha$ -NF), which inhibits CYP1A1 and 1A2; furafylline, which inhibits CYP1A2, diamantane, which inhibits CYP2B (Stiborova et al., 2002a); sulphafenazole, which inhibits CYP2C; diethyldithiocarbamic acid (DDTC), which inhibits CYP2E1 and ketoconazole, which inhibits CYP3A (Rendic & DiCarlo, 1997). Inhibitors were dissolved in ethanol, except of  $\alpha$ -NF that was dissolved in a mixture of methanol:ethylacetate (3:2, v/v) and DDTC that was dissolved in distilled in water, to yield final concentrations of 1-1000 µM in the incubation mixtures.

**HPLC.** HPLC was performed with a reversed phase column (Nucleosil 100-5 C<sub>18</sub>, Macherey-Nagel, Duren, Germany, 25 cm x 4.6 mm, 5 mm) preceded by a C-18 guard column, using isocratic elution conditions of 70% methanol in distilled water with a flow rate of 0.6 ml/min. HPLC was carried out with a Dionex HPLC pump P580 with UV/VIS UVD 170S/340S spectrophotometer detector set at 254 nm, and peaks were integrated with a CHROMELEON™ 6.01 integrator. 3-ABA, *N*-hydroxy-3-ABA, and 3-NBA were eluted with retention times (r.t.) of 8.2, 6.5 and 25 minutes, respectively (Figure 2)

## RESULTS

When 3-ABA was incubated with rat hepatic microsomes in the presence of NADPH up to three product peaks were separated by HPLC (see Figure 2 for hepatic microsomes of rats treated with  $\beta$ -NF). Using co-chromatography with synthetic standards, two of them were identified to be the products formed by oxidation of 3-ABA on its amino group back to the parent compound from which 3-ABA is generated in organisms, 3-NBA. Namely, *N*-hydroxylation metabolite, *N*-hydroxy-3-ABA and 3-NBA were identified to be these 3-ABA oxidation products. This finding indicates that 3-ABA might, in some cases, be oxidized through *N*-hydroxy-3-ABA back to its oxidative counterpart, 3-NBA (Figure 1). A structure of another metabolite eluted with the retention time (r.t.) of 18 min, M18 (Figure 2A) remains to be characterized.

We have used hepatic microsomes of rats treated with CYP inducers (an inducer of CYP1A1/2,  $\beta$ -NF and an inducer CYP2B1/2, PB), as well as inhibitors of individual CYPs, to evaluate the role of rat hepatic CYPs in 3-ABA metabolism (Mizerovska et al., 2008). Here, the same microsomal systems and hepatic microsomes of control (untreated) rats were utilized to determine the kinetics of 3-ABA metabolism. If 3-ABA was incubated with these microsomes more than 30 minutes, different patterns of 3-ABA metabolic products were found in individual microsomes. Whereas hepatic microsomes rich in CYP1A1/2 ( $\beta$ -NF-microsomes) generated the final oxidation metabolite of 3-ABA, 3-NBA (Figure 1), this metabolite (3-NBA) was not formed by the other microsomes tested in the study (Figure 3). The metabolite with unknown structure, M18, was the only one formed from 3-ABA by these enzymatic systems (Figure 3). It should be mentioned that in some cases the control incubations without presence of microsomes contained traces of 3-ABA metabolites, 3-NBA and a metabolite M18 (see Figure 3A showing this case), which might probably be formed

by autooxidation. *N*-hydroxy-3-ABA was another metabolite generated by microsomes rich in CYP1A1/2 (Figure 2A), but this metabolite was not quantified in this study, namely, because this reactive compound is easily decomposed and forms nitrenium and/or carbenium ion (Figure 1), being scavenged by proteins (at least partially) present in the incubation mixtures. When shorter incubation times were used in the experiments, formation of all 3-ABA oxidation products detected in the above experiments was low, being hardly to be quantified. Hence, because the incubation times during them the conversion of 3-ABA is linear (that are shorter than 30 minutes, see below) are needed to study kinetics of 3-ABA metabolism by microsomes, the consumption of 3-ABA, detected during such short incubation times, has been used to evaluate initial velocities of the enzymatic reactions. Conversion of 3-ABA by all three hepatic microsomes (measured by consumption of 3-ABA) was linear up to 5 minutes of incubation (data not shown). This time interval was, therefore, used to evaluate kinetics of 3-ABA oxidation.

A classical Michaelis-Menten kinetics was found for 3-ABA metabolism by hepatic microsomes of rats treated with  $\beta$ -NF and PB; double reciprocal plots of initial velocities *versus* concentrations of 3-ABA were linear (Figure 4A,B). The values of Michaelis constant ( $K_m$ ) and maximum reaction velocity ( $V_{max}$ ) determined from this kinetics are shown in Table 1. The values of  $K_m$  and  $V_{max}$  for 3-ABA metabolism by PB-microsomes were more than 1.2-fold higher than those by hepatic microsomes of rats treated with  $\beta$ -NF.

In contrast to the kinetics found with rat hepatic  $\beta$ -NF- and PB-microsomes, double reciprocal plots of kinetics of 3-ABA oxidation by hepatic microsomes of control (untreated) rats were not linear (Figure 4C), giving two  $K_m$  values for 3-ABA (Table 1). These findings indicate that more than one of the CYP enzymes present in

these microsomes are responsible for 3-ABA metabolism. Because one of the  $K_m$  values is close to the value of  $K_m$  found with microsomes rich in CYP1A1/2 ( $\beta$ -NF-microsomes), the enzyme of a CYP1A subfamily might be important for 3-ABA conversion in hepatic microsomes of control rats. Indeed, inhibitors of CYPs of a 1A subfamily ( $\alpha$ -NF for CYP1A1/2 and furafylline for CYP1A2) are strong inhibitors of 3-ABA metabolism (Table 2). Results of experiments utilizing CYP inhibitors also suggest that CYP enzymes of a 3A subfamily might participate in 3-ABA metabolism in rat hepatic microsomes; a selective CYP3A inhibitor, ketoconazole, was highly effective in inhibiting 3-ABA conversion, with the  $IC_{50}$  value of 3.5  $\mu$ M (Table 2).

To further characterize the role of individual rat CYPs in 3-ABA metabolism and to compare their efficiencies with those of human CYP enzymes, microsomes of Baculovirus transfected insect cells (Supersomes<sup>TM</sup>) containing recombinantly expressed rat and human CYPs and NADPH:CYP reductase were utilized in an additional part of this study.

First, a time-dependence of 3-ABA conversion catalyzed by rat recombinant CYP1A1, 3A1 and 3A2, the enzymes found in the above experiments to be important in 3-ABA metabolism in rat liver microsomes, and that of their human orthologous CYPs (CYP1A1 and 3A4) was investigated. Whereas 3-ABA conversion catalyzed by human and rat CYP1A1, rat CYP3A1 and human CYP3A4 is linear up to 20 minutes of incubations, that by rat CYP3A2 is linear even up to 40 minutes of incubation (data not shown). Because of linearity of 3-ABA conversion till 20 minutes, this time interval was used in experiments evaluating the efficiency of individual human and rat CYPs to metabolize 3-ABA.

All rat and human recombinant CYP tested in this study were effective to metabolize 3-ABA (Figure 5). Among them CYPs of a 3A subfamily (CYP3A2 and

3A4/5) followed by CYP1A1 were the most efficient enzymes of both organisms metabolizing 3-ABA. In addition, rat CYP2C11 and human CYP2A6, 2C19 and 2D6 were also effective in this reaction (Figure 5)

## DISCUSSION

3-ABA, the human metabolite of the ubiquitous environmental pollutant 3-NBA, was detected in the urine of smoking and nonsmoking salt mining workers occupationally exposed to diesel emissions at similar concentration (1-143 ng/24 h urine) to 1-aminopyrene (2-200 ng/24 h urine), the corresponding amine of the most abundant nitro-polycyclic aromatic hydrocarbons detected in diesel exhaust matter (Seidel et al. 2002). Comparison between experimental animals and human CYP enzymes is essential for the extrapolation of animal toxicity data to assess human health risk. Therefore, here we compared the ability of rat and human CYP enzymes to metabolize 3-ABA.

The results found in the presented study have increased our knowledge on the potential of human and rat CYP enzymes to metabolize 3-ABA and on the kinetics of such reactions. Two experimental approaches were utilized in this study. For the first, we used hepatic microsomal systems of rat found to be a suitable experimental model (based on the fact that the same enzymes activate 3-ABA in human and rat livers to form DNA adducts) (Arlt et al. 2004b; 2006c, Stiborova et al. 2006a). Therefore, the results should provide some indications of what might occurs with 3-ABA in livers of human. The second approach employed rat and human recombinant CYP enzymes, which should mimic the enzymatic activity of actual CYP enzymes of both organisms.

297        Recently, we have found that rat and human hepatic CYP1A1 and 1A2,  
298 followed by human CYP2A6, are predominantly responsible for metabolic activation  
299 of 3-ABA to form DNA adducts. The extrahepatic enzymes CYP1B1 and 2B6 were  
300 also effective to activate 3-ABA, but to a much lower extent (Arlt et al. 2004b). On the  
301 contrary, none of the other CYP enzymes such as CYP2C9, 2D6, 2E1 and 3A4  
302 tested in this former study activated 3-ABA to species generating DNA adducts (Arlt  
303 et al. 2004b). This finding corresponds to results found in the present study. We have  
304 found that CYP1A enzymes are effective in oxidation of 3-ABA to the final oxidative  
305 metabolite of this substance, 3-NBA. Namely, to the metabolite that is formed  
306 through the formation of *N*-hydroxy-3-ABA. This reactive intermediate (*N*-hydroxy-3-  
307 ABA) found in this study to be formed by CYP1A enzymes, is also decomposed to  
308 the ultimate carcinogenic species of 3-ABA, nitrenium and/or carbenium ions, forming  
309 the 3-ABA-derived DNA adducts (Figure 1). Hence, this metabolic activation pathway  
310 seems to be mediated mainly by the CYP1A enzymes. On the contrary, 3-ABA  
311 oxidation to further product(s) such as metabolite M18, which is supposed to be a  
312 detoxification metabolite (Mizerovska et al. 2008), is catalyzed not only by CYP1A,  
313 but also by other rat CYP enzymes. As follows from the studies on kinetics of 3-ABA  
314 metabolism (presented study) as well as from the effects of selective inhibitors of  
315 individual CYPs (Mizerovska et al. 2008 and presented study), beside CYP1A, the  
316 CYP enzymes of 3A and 2C subfamilies might be very effective in such 3-ABA  
317 metabolism in rat livers. Using rat recombinant CYPs, this suggestion was confirmed;  
318 rat CYP3A2 followed by CYP1A1, 2C11 and 3A1 were the most efficient in 3-ABA  
319 conversion. The results of this work also support our former findings showing a  
320 suitability of the rat as an appropriate animal model to study the fate of 3-ABA in  
321 humans (Arlt et al. 2004b; Stiborova et al. 2006a). Namely, here we have found that

human CYPs of a 3A subfamily (CYP3A and 3A5), orthologous CYPs to rat enzymes mentioned above, followed by human and rat CYP1A1, are the most efficient in 3-ABA metabolism.

In conclusion, the present study characterizing two metabolites formed from 3-ABA by CYP-mediated oxidation, namely, a metabolite of 3-ABA, which is responsible for generation of 3-ABA-derived DNA adducts (N-hydroxy-3-ABA) and its final oxidation product (3-NBA), confirmed the participation of CYP1A enzymes in metabolic activation of 3-ABA (Arlt et al. 2004b). Other rat and human CYP enzymes, such a CYP3A and 2C are mainly important for detoxification metabolism of this compound. Structural characterization of the major detoxification metabolite of 3-ABA awaits further investigation.

**ACKNOWLEDGMENT** Supported by the Grant Agency of the Czech Republic (grants 303/09/0472, 203/09/0812 and 305/09/H008) and the Ministry of Education of the Czech Republic (grants MSM0021620808 and 1M0808).

## REFERENCES

- 1 Arlt VM (2005). 3-Nitrobenzanthrone, a potential human cancer hazard in diesel exhaust and urban air pollution: A review of the evidence. *Mutagenesis*. **20**: 399–410.
- 2 Arlt VM, Bieler CA, Mier W, Wiessler M and Schmeiser HH (2001). DNA adduct formation by the ubiquitous environmental contaminant 3-nitrobenzanthrone in rats determined by <sup>32</sup>P-postlabeling. *Int J Cancer*. **93**: 450–454.

- 346 3 Arlt VM, Glatt HR, Muckel E, Pabel U, Sorg BL, Schmeiser HH and Phillips DH  
347 (2002). Metabolic activation of the environmental contaminant 3-  
348 nitrobenzanthrone by human acetyltransferases and sulfotransferase.  
349 Carcinogenesis. **23**: 1937–1945.
- 350 4 Arlt VM, Glatt H, Muckel E, Pabel U, Sorg BL, Seidel A, Frank H, Schmeiser  
351 HH and Phillips DH (2003a). Activation of 3-nitrobenzanthrone and its  
352 metabolites by human acetyltransferases, sulfotransferases and cytochrome  
353 P450 expressed in Chinese hamster V79 cells. Int J Cancer. **105**: 583–592.
- 354 5 Arlt VM, Sorg BL, Osborne M, Hewer A, Seidel A, Schmeiser HH and Phillips  
355 DH (2003b). DNA adduct formation by the ubiquitous environmental pollutant  
356 3-nitrobenzanthrone and its metabolites in rats. Biochem Biophys Res  
357 Commun. **300**: 107–114.
- 358 6 Arlt VM, Stiborova M, Hewer A, Schmeiser HH and Phillips DH, (2003c).  
359 Human enzymes involved in the metabolic activation of the environmental  
360 contaminant 3-nitrobenzanthrone: evidence for reductive activation by human  
361 NADPH:cytochrome P450 reductase. Cancer Res. **63**: 2752-2761.
- 362 7 Arlt VM, Cole KJ and Phillips DH (2004a). Activation of 3-nitrobenzanthrone  
363 and its metabolites to DNA-damaging species in human B-lymphoblastoid  
364 MCL-5 cells. Mutagenesis. **19**: 149–156.
- 365 8 Arlt VM, Hewer A, Sorg BL, Schmeiser HH, Phillips DH and Stiborova M  
366 (2004b). 3-Aminobenzanthrone, a human metabolite of the environmental  
367 pollutant 3-nitrobenzanthrone, forms DNA adducts after metabolic activation  
368 by human and rat liver microsomes: evidence for activation by cytochrome  
369 P450 1A1 and P450 1A2. Chem Res Toxicol. **17**: 1092–1101.



- 370 9 Arlt, VM, Zhan L, Schmeiser HH, Honma M, Hayashi M, Phillips DH and  
371 Suzuki T (2004c). DNA adducts and mutagenic specificity of the ubiquitous  
372 environmental pollutant 3-nitrobenzanthrone in Muta Mouse. *Environ Mol*  
373 *Mutagen.* **43**: 186–195.
- 374 10 Arlt VM, Stiborova M, Henderson CJ, Osborne MR, Bieler CA, Frei E, Martinek  
375 V, Sopko B, Wolf CR, Schmeiser HH and Phillips DH (2005). The  
376 environmental pollutant and potent mutagen 3-nitrobenzanthrone forms DNA  
377 adducts after reduction by NAD(P)H:quinone oxidoreductase and conjugation  
378 by acetyltransferases and sulfotransferases in human hepatic cytosols.  
379 *Cancer Res.* **65**: 2644–2652.
- 380 11 Arlt VM, Henderson CJ, Wolf CR, Schmeiser HH, Phillips DH and Stiborova M  
381 (2006a). Bioactivation of 3-aminobenzanthrone, a human metabolite of the  
382 environmental pollutant 3-nitrobenzanthrone: evidence for DNA adduct  
383 formation mediated by cytochrome P450 enzymes and peroxidases. *Cancer*  
384 *Lett.* **234**: 220–231.
- 385 12 Arlt VM, Schmeiser HH, Osborne MR, Kawanishi M, Kanno T, Yagi T, Phillips  
386 DH and Takamura-Enya T (2006b). Identification of three major DNA adducts  
387 formed by the carcinogenic air pollutant 3-nitrobenzanthrone in rat lung at the  
388 C8 and N<sup>2</sup> position of guanine and at the N<sup>6</sup> position of adenine. *Int J Cancer.*  
389 **118**: 2139–2146.
- 390 13 Bieler CA, Wiessler M, Erdinger L, Suzuki H, Enya T and Schmeiser HH  
391 (1999). DNA adduct formation from the mutagenic air pollutant 3-  
392 nitrobenzanthrone. *Mutat Res.* **439**: 307–311.
- 393 14 Bieler CA, Cornelius M, Klein R, Arlt VM, Wiessler M, Phillips DH and  
394 Schmeiser HH (2005). DNA adduct formation by the environmental

395 contaminant 3-nitrobenzanthrone after intratracheal instillation in rats. *Int J*  
396 *Cancer*. **118**: 833–838.

397 15 Bieler CA, Cornelius MG, Stiborova M, Arlt VM, Wiessler M, Phillips DH and  
398 Schmeiser HH (2007). Formation and persistence of DNA adducts formed by  
399 the carcinogenic air pollutant 3-nitrobenzanthrone in target and non-target  
400 organs after intratracheal instillation in rats. *Carcinogenesis*. **28**: 1117–1121.

401 16 Borlak J, Hansen T, Yuan Z, Sikka HC, Kumar S, Schmidbauer S, Frank H,  
402 Jacob J and Seidel A (2000). Metabolism and DNA-binding of 3-  
403 nitrobenzanthrone in primary rat alveolar type II cells, in human fetal bronchial,  
404 rat epithelial and mesenchymal cell lines. *Polycyclic Aromat Compounds*. **21**:  
405 73–86.

406 17 Enya T, Suzuki H, Watanabe T, Hirayama T and Hisamatsu Y (1997). 3-  
407 Nitrobenzanthrone, a powerful bacterial mutagen and suspected human  
408 carcinogen found in diesel exhausts and airborne particulates. *Environ Sci*  
409 *Technol*. **31**: 2772–2776.

410 18 Grabchev I, Moneva I, Betcheva R and Elyashevich G (2002). Colored  
411 microporous polyethylene films: effect of porous structure on dye adsorption.  
412 *Mat Res Innovat*. **6**: 34–37.

413 19 Hansen T, Seidel A and Borlak J (2007). The environmental carcinogen 3-  
414 nitrobenzanthrone and its main metabolite 3-aminobenzanthrone enhance  
415 formation of reactive oxygen intermediates in human A549 lung epithelial  
416 cells. *Toxicol Appl Pharmacol*. **221**: 222–234.

417 20 Krizkova J, Burdova K, Hudecek J, Stiborova M, Hodek P (2008). Induction of  
418 cytochromes P450 in small intestine by chemopreventive compounds.  
419 *Neuroendocrinol. Lett*. **29**, 717-721.

- 420 21 Mizerovska J, Dracinska H, Arlt VM, Hudecek J, Hodek P, Schmeiser HH, Frei  
421 E, Stiborova M (2008). Rat cytochromes P450 oxidize 3-aminobenzanthrone,  
422 a human metabolite of the carcinogenic environmental pollutant 3-  
423 nitrobenzanthrone. *Interdisc Toxicol.* **1**: 150–154
- 424 22 Nagy E, Zeisig M, Kawamura K, Hisumatsu Y, Sugeta A, Adachi S and Moller  
425 L (2005). DNA-adduct and tumor formations in rats after intratracheal  
426 administration of the urban air pollutant 3-nitrobenzanthrone. *Carcinogenesis.*  
427 **26**: 1821–1828.
- 428 23 Omura T, Sato R (1964). The carbon monoxide-binding pigment of liver  
429 microsomes. I. Evidence for its hemoprotein nature. *J Biol Chem.* **239**: 2370–  
430 2378
- 431 24 Rendic S, DiCarlo FJ (1997). Human cytochrome P450 enzymes: A status  
432 reportsummarizing their reactions, substrates, inducers, and inhibitors. *Drug*  
433 *Metab. Rev.* **29**: 413–80.
- 434 25 Seidel A, Dahmann D, Krekeler H and Jacob J (2002). Biomonitoring of  
435 polycyclic aromatic compounds in the urine of mining workers occupationally  
436 exposed to diesel exhaust. *Int J Hyg Environ Health.* **204**: 333–338.
- 437 26 Sistkova J, Hudecek J, Hodek P, Frei E, Schmeiser HH and Stiborova M  
438 (2008). Human cytochromes P450 1A1 and 1A2 participate in detoxication of  
439 carcinogenic aristolochic acid. *Neuroendocrinol. Lett.* **29**, 733–737.
- 440 27 Stiborova M., Borek-Dohalska L., Hodek P., Mraz J., Frei E. (2002a). New  
441 selective inhibitors of cytochromes P450 2B and their application to  
442 antimutagenesis of tamoxifen. *Arch. Biochem. Biophys.* **403**: 41–49.
- 443 28 Stiborova M, Martinek V, Rydlova H, Hodek P and Frei E (2002b). Sudan I is a  
444 potential carcinogen for humans: evidence for its metabolic activation and

445 detoxication by human recombinant cytochrome P450 1A1 and liver  
 446 microsomes. *Cancer Res.* **62**: 5678–5684.

447 29 Stiborova M, Dracinska H, Hajkova J, Kaderabkova P, Frei E, Schmeiser HH,  
 448 Soucek P, Phillips DH and Arlt VM (2006a). The environmental pollutant and  
 449 carcinogen 3-nitrobenzanthrone and its human metabolite 3-  
 450 aminobenzanthrone are potent inducers of rat hepatic cytochromes P450 1A1  
 451 and -1A2 and NAD(P)H:quinone oxidoreductase. *Drug Metab Dispos* **34**:  
 452 1398–1405.

453 30 Stiborova M., Martinek V., Schmeiser H.H., Frei E. (2006b). Modulation of  
 454 CYP1A1-mediated oxidation of carcinogenic azo dye Sudan I and its binding  
 455 to DNA by cytochrome b<sub>5</sub>. *Neuroendocrinol. Lett.* **27** (Suppl. 2), 35–39.

456 31 Stiborova M, Dracinska H, Mizerovska J, Frei E, Schmeiser HH, Hudecek J,  
 457 Hodek P, Phillips DH and Arlt VM (2008). The environmental pollutant and  
 458 carcinogen 3-nitrobenzanthrone induces cytochrome P450 1A1 and  
 459 NAD(P)H:quinone oxidoreductase in rat lung and kidney, thereby enhancing  
 460 its own genotoxicity. *Toxicology.* **247**: 11–22.

461 32 Stiborova M., Dracinska H., Martinkova M., Mizerovska J., Hudecek J., Hodek  
 462 P., Liberda J., Frei E., Schmeiser H.H., Phillips D.H., Arlt V.M (2009). 3-  
 463 Aminobenzanthrone, a human metabolite of the carcinogenic environmental  
 464 pollutant 3-nitrobenzanthrone, induces biotransformation enzymes in rat  
 465 kidney and lung. *Mutat. Res.- Gen. Tox. En.* **676**: 93–101.

466 33 Svobodova M, Sistkova J, Dracinska H, Hudecek J, Hodek P, Schmeiser HH,  
 467 Arlt VM, Frei E and Stiborova M (2007). Reductive activation of environmental  
 468 pollutants 3-nitrobenzanthrone and 2-nitrobenzanthrone. *Chem Listy.* **100**:  
 469 s277–s279

- 34 Sykora J, Mudago V, Hutterer R, Nepras M, Vanerka J, Kapusta P, Fidler V  
and Hof M (2002). ABA-C-15: A new dye for probing solvent relaxation in  
phospholipid bilayers. *Langmuir*. **18**: 9276–9282.
- 35 Weichelman KJ, Braun RD, Fitzpatrick JD (1988). Investigation of the  
bicinchoninic acid protein assay: identification of the groups responsible for  
color formation. *Anal Biochem*. **175**: 231–237

Table 1. Kinetic parameters of 3-ABA oxidation catalyzed by hepatic microsomal CYPs

Parameters			
Hepatic microsomes from rats pretreated with	$K_m$		$V_{max}$
	$K_{m1}$	$K_{m2}$	$V_{max1}$ $V_{max2}$
	[ $\mu M$ ]		[nmol 3-ABA min <sup>-1</sup> mg <sup>-1</sup> ]
$\beta$ -naphthoflavone (CYP1A1/2)	47.5 $\pm$ 0.5		13.42 $\pm$ 0.134
Phenobarbital (CYP2B)	60.7 $\pm$ 0.6		16.8 $\pm$ 0.168
Control	20.7 $\pm$ 0.2	417.5 $\pm$ 4.18	3.4 $\pm$ 0,3      48.5 $\pm$ 4.85

Experimental conditions are described in Material and methods, 3-ABA (5 - 50  $\mu M$ ) and 0.5 mg microsomal protein were present in the incubation mixtures to determine the  $K_m$  and  $V_{max}$  values. Values in the table are averages and standard deviations of three determinations.

Table 2. The effects of CYP inhibitors on 3-ABA oxidation by rat hepatic microsomes

Hepatic microsomes from rats pretreated with <sup>a</sup>	Inhibitor <sup>b</sup>	IC <sub>50</sub> [μM] <sup>c</sup>
β-naphtoflavone (CYP1A1/2)	α-naphtoflavone (CYP1A1/2)	10.9 ± 1.0
β-naphtoflavone (CYP1A1/2)	Furafylline (CYP1A2)	16.6 ± 1.6
Untreated	Sulfaphenazole (CYP2C)	10.8 ± 1.0
Untreated	DDTC (CYP2E1)	96.8 ± 6.1
Untreated	Ketoconazole (CYP3A4)	3.5 ± 0.3

<sup>a</sup>Isoforms of CYPs induced by inducers are shown in brackets. <sup>b</sup>Isoforms of CYP inhibited by selective inhibitors are shown in brackets. <sup>c</sup>Estimated from concentration-dependent inhibition of 3-ABA oxidation by interpolation (inhibitors were 1 – 1000 μM depending on the chemical). 3-ABA (20 μM) and 0.5 mg microsomal protein were present in the incubation mixture. <sup>d</sup>Averages and standard deviations of three determinations.

Figure 1. Pathways of metabolism and DNA adduct formation of 3-NBA and 3-ABA. See text for details. NQO1, NAD(P)H:quinone oxidoreductase; NAT, *N,O*-acetyltransferases; SULT, sulfotransferase; PHS-1, prostaglandin H synthase-1 (cyclooxygenase 1); CYP, cytochrome P450; LPO, lactoperoxidase; MPO, myeloperoxidase; POR, NADPH:cytochrome P450 oxidoreductase; R = -COCH<sub>3</sub> or -SO<sub>3</sub>H. R = -COCH<sub>3</sub> or -SO<sub>3</sub>H; dA-*N*<sup>6</sup>-ABA, 2-(2'-deoxyadenosin-*N*<sup>6</sup>-yl)-3-aminobenzanthrone; dG-*N*<sup>2</sup>-ABA, *N*-(2'-deoxyguanosin-*N*<sup>2</sup>-yl)-3-aminobenzanthrone; dG-C8-*N*-ABA, *N*-(2'-deoxyguanosin-8-yl)-3-aminobenzanthrone.






Figure 2. HPLC chromatographs of 3-ABA metabolites produced by hepatic microsomes of rats treated with  $\beta$ -NF (A), HPLC profiles of several standards; 3-ABA (B), *N*-hydroxy-3-ABA (C) and 3-NBA (D).




Figure 3. Time dependence of 3-ABA oxidation by hepatic microsomes of rats treated with  $\beta$ -NF (A), PB (B), and by those of control (uninduced) rats (C). Experimental conditions are described in Material and methods. Values of reaction rates of 3-ABA oxidation are averages and standard deviations of triplicate incubation. MS; microsomes




Figure 4. Double reciprocal plots of initial velocities of 3-ABA oxidation *versus* concentrations of 3-ABA catalyzed by hepatic microsomes of rats treated with PB (A),  $\beta$ -NF (B) and those of control (untreated) rats (C). Reaction rate ( $v$ ) [nmol 3-ABA/min/mg protein]. Experimental conditions are described in Material and methods.




Figure 5. Oxidation of 3-ABA by recombinant CYPs of rat (A) and human (B). Experimental conditions are described in Material and methods. Values of reaction rates of 3-ABA oxidation are averages and standard deviations of triplicate incubation.

